

UWMC ED - Anticoagulant Reversal for SAH & IPH:

Warfarin?

- Give Vitamin K 10 mg IV over one hour (empiric reversal)
- Request 4 units of universal-donor FFP from UWMC Transfusion Support Services (x8-8700) to bedside while awaiting blood test results
- If INR ≥ 1.5 , transfuse 2 units universal-donor (AB) thawed plasma.
- Refer to [Guidelines for Reversing Coagulopathies](#)
 - Discuss with Neurology &/ or Stroke Attending whether to use Bebulin pathway or FFP pathway

Antiplatelet therapy?

- For aspirin, clopidogrel, prasugrel, NSAIDs, glycoprotein IIb/IIIa antagonists (abciximab, eptifibatide, or tirofiban):*
- Request platelets from Puget Sound Blood Center (206. 522-2462) while awaiting blood test results
 - Ask for 2 U universal-donor apheresed platelets
 - Give the platelets as soon as possible irregardless of the lab test results

Heparinoids?

- If on **full-dose unfractionated heparin**, stop it and give:
 - Give Protamine 25 mg IV before blood results return
 - After infusion, repeat STAT PTT
 - Refer to [Guidelines for Reversing Coagulopathies](#)
- If on **low-molecular-weight heparin** (*enoxaparin, ardeparin, certoparin, dalteparin, nadroparin, reviparin, and tinzaparin*) within:
 - The last 24 hours:
 - Give Protamine 25 mg IV as soon as possible
 - The last 8 hours:
 - Give Protamine 50 mg IV as soon as possible
- Refer to [Guidelines for Reversing Coagulopathies](#)

Unknown therapy?

- *If on some other type of unfamiliar antithrombotic treatment, consult hematology through the page operator*